

AQA C2 Bonding and Structure

Triple Chemistry Page 1 of 2


Bonding occurs because chemicals are only stable when the particles have full outer shells of electrons

Keywords

atom	the smallest particle of a chemical element that can exist
element	a chemical made up of only one type of atom
ion	a particle which has a positive or negative charge (lost or gained electrons)
electrostatic force	the attraction between positively and negatively charged particles
bond	strong attraction between atoms or ions
state (of matter)	whether a substance is a solid, liquid or gas
molecule	a small group of atoms held together by covalent bonds
alloy	a material which contains a metal and at least one other element
delocalised	free to move
malleable	can be bent and shaped
molten	liquid
intermolecular	forces between molecules

Alloys

Alloys contain a mixture of a metal and at least one other element. They are harder than pure metals. This is because the layers of ions can't slide over each other due to the different sizes.



States of matter			Limitations of model: particles are shown as solid spheres and there are no forces between the particles
state	model	state symbol	
solid		(s)	
liquid		(l)	
gas		(g)	

Electrical conductivity

For a material to conduct electricity it needs to have:

- charged particles (electrons or ions)
- which can move

Metallic bonding – seen in metals and alloys

electrons	Electrons in the outer shells of metals are delocalised forming positive metal ions	
structure	Metallic structure held together by strong electrostatic forces between the lattice of positive ions and the delocalised electrons	
properties	<p>High melting / boiling points (a lot of energy needed to overcome strong metallic bonds). Conduct electricity (delocalised electrons carry charge through the metal). Conduct thermal energy (delocalised electrons move through the structure transferring energy). Malleable (layers of ions slide over each other)</p>	






Changes of state

melting point	At this temperature: <ul style="list-style-type: none"> solids melt liquids freeze 	Stronger attraction: <ul style="list-style-type: none"> more energy to overcome bond higher melting / boiling point
boiling point	At this temperature: <ul style="list-style-type: none"> liquids boil gases condense 	Weaker attraction: <ul style="list-style-type: none"> less energy to overcome bond lower melting / boiling point


Forming ions	Metals lose electrons forming positive ions	Non-metals gain electrons forming negative ions



Ionic bonding – between a metal and a non-metal



electrons	Electrons are transferred from the metal to the non-metal forming ions	Dot and cross diagram
structure	Giant ionic lattice held together by strong electrostatic forces between positive and negative ions	
properties	<p>High melting / boiling points (a lot of energy is needed to overcome strong ionic bonds). When solid they do not conduct electricity (ions are held in fixed positions within a lattice and cannot move). When dissolved or molten they do conduct electricity (when the lattice breaks apart, the ions are free to move and carry charge).</p>	


Structure and bonding of carbon	
diamond	 <p>Each carbon atom forms four covalent bonds with other carbon atoms in a giant covalent structure. Because covalent bonds are strong diamond is very hard and has a very high melting point. It does not conduct electricity as the electrons are held between the atoms.</p>
graphite	 <p>Each carbon atom forms three covalent bonds with three other carbon atoms, forming layers. There are weak forces between the layers so they can easily slide over each other.</p>
graphene	 <p>Graphene is a single layer of graphite. It has a high melting and boiling point and can conduct electricity, making it useful in electronics and composites.</p>
carbon nanotubes	 <p>Carbon nanotubes are cylindrical fullerenes. They are used for electronics, nanotechnology and materials.</p>
fullerenes	 <p>Fullerenes are large molecules of carbon atoms with hollow shapes. The first to be discovered was Buckminsterfullerene (C₆₀). This looks like a sphere.</p>


Three electrons from the outer shell of each carbon atom form covalent bonds, then the fourth electron is delocalised. Therefore these structures can conduct electricity.

Giant covalent structures – bonds between non-metal atoms	
electrons	All of the atoms are linked to other atoms by strong covalent bonds forming a giant covalent structure . Examples are diamond, graphite and silicon dioxide
structure	<p>Silicon dioxide</p>  <p>oxygen atom, silicon atom, covalent bond</p>
properties	High melting / boiling points (a lot of energy is needed to overcome strong covalent bonds). Do not conduct electricity (electrons are in bonds so cannot move or carry charge).

Polymers	
structure	<p>These are very large molecules containing atoms linked to other atoms by strong covalent bonds.</p>  <p>poly(ethene)</p>
properties	<p>Normally solids at room temperature (the forces between the molecules are fairly strong).</p>  <p>fairly strong forces of attraction between the polymers</p>

Covalent molecular structures – bonding between non-metals	
electrons	<p>Atoms share pairs of electrons forming strong covalent bonds between the atoms.</p> <p>Dot-cross diagram (eg. ammonia – NH₃):</p> 
structure	<p>Small molecules which have <u>strong covalent bonds</u> but <u>weak intermolecular forces</u> (forces between molecules)</p>  <p>nitrogen molecule, Intermolecular force, Intramolecular force</p>
properties	<p>Usually gases or liquids (low melting and boiling points) Low melting and boiling points (weak intermolecular forces don't need much energy to overcome). Melting and boiling points increase as molecules get bigger (intermolecular forces are stronger when molecules are bigger). Do not conduct electricity (molecules are neutral so there are no charged particles).</p>

Nanoparticles	
Size:	<p>1-100 nm (1 nm = 10⁻⁹ m)</p> <p>Each particle contains a few hundred atoms</p>
Surface area : volume ratio (SA : V)	<p>This ratio is very large in nanoparticles and can change the properties of the substance a lot</p> <p>In solids: smaller particles = larger SA : V</p>  <p>SA : V = 3 : 5 SA : V = 6 : 1</p>
Uses	<ul style="list-style-type: none"> • medicine • electronics • cosmetics • sun creams • deodorants • catalysts

Models
<p>Dot-cross, ball and stick (a), 2D and 3D (b) models all have limitations such as not showing the bonding electrons, not showing the chemical bonds, or not showing the 3D structure of materials.</p>  <p>(a) (b)</p>