



Alkenes




Ethene C₂H₄




Propene C₃H₆

alkenes	unsaturated hydrocarbons
unsaturated	this means they contain a double bond
general formula	C _n H _{2n}


functional group	Alkenes are hydrocarbons in the functional group C=C	The functional group of an organic compound determines their reaction
alkene reactions	Alkenes react with oxygen in the same way as other hydrocarbons, just with a smoky flame due to incomplete combustion	Alkenes also react with hydrogen, water and the halogens. The C=C bond allows for the addition of other atoms. The pictures below show some examples.




+



chlorine, bromine or iodine can be added




↓



alkene + water (steam) → alcohol

This is called hydration, and it needs a temperature of approximately 300°C and an acid catalyst.




alkene + hydrogen → alkane
This is called hydrogenation, and it needs a nickel catalyst.


All of these are examples of reactions where molecules add on each side of the double bond, and the double bond breaks to leave a single bond.

Alcohols


Functional group	Alcohol reactions	Fermentation
-OH For example: CH ₃ CH ₂ OH	Alcohols react with sodium, air and water. They are useful as fuels and solvents	Ethanol is produced from fermentation
Methanol, ethanol, propanol and butanol are the first four of the homologous series.	Alcohols and sodium: bubbling, hydrogen gas given off and salt formed Alcohols and air: alcohols burn in air releasing carbon dioxide and water Alcohols and water: alcohols dissolve in water to form a neutral solution	When sugar solutions are fermented using yeast, aqueous solutions of ethanol are produced. The conditions needed for this process include a moderate temperature (25-50°C), water (from sugar solution) and an absence of oxygen.




Methanol



Ethanol





Propanol



Butanol

Carboxylic acids

Functional group	Carboxylic acid reactions	Strength (HT only)
-COOH For example: CH ₃ COOH	Carboxylic acids react with carbonates (to fizz and form CO ₂). For example, ethanoic acid + sodium carbonate → sodium ethanoate + water + carbon dioxide	Carboxylic acids are weak acids. They only partially ionise in water. An aqueous solution of a weak acid will have a high pH (but still below 7).
Methanoic acid, ethanoic acid, propanoic acid and butanoic acid are the first four of the homologous series.	Carboxylic acids dissolve in water. Carboxylic acids react with alcohols to form esters.	Carboxylic acids react with alcohols to make <u>esters</u> . Esters are volatile and have characteristic smells



Ethyl ethanoate