


**Bonding occurs because chemicals are only stable when the particles have full outer shells of electrons**

**Keywords**

<b>atom</b>	the smallest particle of a chemical element that can exist
<b>element</b>	a chemical made up of only one type of atom
<b>ion</b>	a particle which has a positive or negative charge
<b>electrostatic force</b>	the attraction between positively and negatively charged particles
<b>(chemical) bond</b>	the force of attraction that holds particles together
<b>state (of matter)</b>	whether a substance is a solid, liquid or gas
<b>molecule</b>	a small group of atoms held together by covalent bonds
<b>alloy</b>	a material which contains a metal and at least one other element
<b>delocalised</b>	free to move
<b>malleable</b>	can be bent and shaped
<b>molten</b>	liquid
<b>intermolecular</b>	forces between molecules
<b>intramolecular</b>	covalent bonds within molecules

**Alloys**

Alloys contain a mixture of a metal and at least one other element. They have the same properties as metals, except that they are harder than pure metals. This is because the layers of ions can't slide over each other due to the different sizes.



States of matter		
state	model	state symbol
solid		(s)
liquid		(l)
gas		(g)

Changes of state		
<b>melting point</b>	At this temperature: <ul style="list-style-type: none"> <li>solids melt</li> <li>liquids freeze</li> </ul>	<b>Stronger bond:</b> <ul style="list-style-type: none"> <li>more energy needed to overcome bond</li> <li>higher melting / boiling point</li> </ul>
<b>boiling point</b>	At this temperature: <ul style="list-style-type: none"> <li>liquids boil</li> <li>gases condense</li> </ul>	<b>Weaker bond:</b> <ul style="list-style-type: none"> <li>less energy needed to overcome bond</li> <li>lower melting / boiling point</li> </ul>






Electrical conductivity
For a material to conduct electricity it needs to have: <ul style="list-style-type: none"> <li>charged particles (electrons or ions)</li> <li>which can move</li> </ul>

Forming ions	Metals lose electrons forming positive ions	Non-metals gain electrons forming negative ions

Metallic bonding – seen in metals and alloys	
<b>electrons</b>	Electrons in the outer shells of metals are <b>delocalised</b> forming positive metal ions
<b>structure</b>	<p><b>Metallic structure</b> held together by <b>strong electrostatic forces</b> between the lattice of positive ions and the delocalised electrons</p>
<b>properties</b>	<p><b>High melting / boiling points</b> (a lot of energy needed to overcome strong metallic bonds).  <b>Conduct electricity</b> (delocalised electrons carry charge through the metal).  <b>Conduct thermal energy</b> (delocalised electrons move through the structure transferring energy).  <b>Malleable</b> (layers of ions slide over each other)</p>


Ionic bonding – between a metal and a non-metal	
<b>electrons</b>	Electrons are transferred from the metal to the non-metal forming ions
<b>structure</b>	<p><b>Dot and cross diagram</b></p> <p><b>Giant ionic lattice</b> held together by <b>strong electrostatic forces</b> between positive and negative ions</p>
<b>properties</b>	<p><b>High melting / boiling points</b> (a lot of energy is needed to overcome strong ionic bonds).  <b>When solid they do not conduct electricity</b> (ions are held in fixed positions within a lattice and cannot move).  <b>When dissolved or molten they do conduct electricity</b> (when the lattice breaks apart, the ions are free to move and carry charge).</p>

**Structure and bonding of carbon**



<b>diamond</b>		Each carbon atom forms four covalent bonds with other carbon atoms in a giant covalent structure. Because covalent bonds are strong diamond is very hard and has a very high melting point. It does not conduct electricity as the electrons are held between the atoms.
<b>graphite</b>		Each carbon atom forms three covalent bonds with three other carbon atoms, forming layers of hexagonal rings. There are weak forces between the layers so they can easily slide over each other.
<b>graphene</b>		<b>Graphene</b> is a single layer of graphite. It has a high melting and boiling point and can conduct electricity, making it useful in electronics and composites.
<b>carbon nanotubes</b>		<b>Carbon nanotubes</b> are cylindrical fullerenes with very high length to diameter ratios. They are used for electronics, nanotechnology and materials.
<b>fullerenes</b>		<b>Fullerenes</b> are large molecules of carbon atoms with hollow shapes. They contain rings of 5, 6, or 7 carbon atoms. The first to be discovered was Buckminsterfullerene (C <sub>60</sub> ).

Three electrons from the outer shell of each carbon atom form covalent bonds, then the fourth electron is delocalised. Therefore these structures can conduct electricity.



**Giant covalent structures – bonds between non-metal atoms**

<b>electrons</b>	All of the atoms are linked to other atoms by strong covalent bonds forming a <b>giant covalent structure</b> . Examples are <b>diamond, graphite and silicon dioxide</b>
<b>structure</b>	<b>Silicon dioxide</b>  oxygen atom, silicon atom, covalent bond
<b>properties</b>	<b>High melting / boiling points</b> (a lot of energy is needed to overcome strong covalent bonds). <b>Do not conduct electricity</b> (electrons are localised in bonds so cannot move or carry charge).





**polymers**

<b>structure</b>	These are very large molecules containing atoms linked to other atoms by strong covalent bonds.  poly(ethene)
<b>properties</b>	<b>Normally solids at room temperature</b> (the forces between the molecules are fairly strong).  fairly strong forces of attraction between the polymers

**Covalent molecular structures – bonding between non-metals**

<b>electrons</b>	Atoms <b>share pairs of electrons</b> forming strong covalent bonds between the atoms. 
<b>structure</b>	<b>Small molecules</b> which have strong intramolecular covalent bonds (bonds within molecules) but weak intermolecular forces of attraction (forces between molecules) 
<b>properties</b>	<b>Usually gases or liquids</b> (low melting and boiling points) <b>Low melting and boiling points</b> (weak intermolecular forces don't need much energy to overcome). <b>Melting and boiling points increase as molecules get bigger</b> (intermolecular forces are stronger when molecules have a higher mass). <b>Do not conduct electricity</b> (molecules are neutral so there are no charged particles).

**Models of chemicals**

Dot-cross diagrams		Ball and stick	
	<b>+ve:</b> shows the pairs of electrons <b>-ve:</b> doesn't show how the particles are arranged in 3D		<b>+ve:</b> shows the covalent bonds <b>-ve:</b> doesn't show the pairs of electrons
2D diagrams		3D diagrams	
	<b>+ve:</b> shows the ions arranged in a lattice <b>-ve:</b> doesn't show how the particles are arranged in 3D		<b>+ve:</b> shows how the particles are arranged in 3D <b>-ve:</b> doesn't show the bonds between the particles