

# Completing the square

## Key points

- Completing the square for a quadratic rearranges  $ax^2 + bx + c$  into the form  $p(x + q)^2 + r$
- If  $a \neq 1$ , then factorise using  $a$  as a common factor.

## Examples

**Example 1** Complete the square for the quadratic expression  $x^2 + 6x - 2$

$x^2 + 6x - 2$ $= (x + 3)^2 - 9 - 2$ $= (x + 3)^2 - 11$	<p><b>1</b> Write <math>x^2 + bx + c</math> in the form</p> $\left(x + \frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2 + c$ <p><b>2</b> Simplify</p>
---	--

**Example 2** Write  $2x^2 - 5x + 1$  in the form  $p(x + q)^2 + r$

$2x^2 - 5x + 1$ $= 2\left(x^2 - \frac{5}{2}x\right) + 1$ $= 2\left[\left(x - \frac{5}{4}\right)^2 - \left(\frac{5}{4}\right)^2\right] + 1$ $= 2\left(x - \frac{5}{4}\right)^2 - \frac{25}{8} + 1$ $= 2\left(x - \frac{5}{4}\right)^2 - \frac{17}{8}$	<p><b>1</b> Before completing the square write <math>ax^2 + bx + c</math> in the form</p> $a\left(x^2 + \frac{b}{a}x\right) + c$ <p><b>2</b> Now complete the square by writing <math>x^2 - \frac{5}{2}x</math> in the form</p> $\left(x + \frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2$ <p><b>3</b> Expand the square brackets – don't forget to multiply <math>\left(\frac{5}{4}\right)^2</math> by the factor of 2</p> <p><b>4</b> Simplify</p>
--	---

## Practice

- 1 Write the following quadratic expressions in the form  $(x + p)^2 + q$
- |                         |                          |
|-------------------------|--------------------------|
| <b>a</b> $x^2 + 4x + 3$ | <b>b</b> $x^2 - 10x - 3$ |
| <b>c</b> $x^2 - 8x$     | <b>d</b> $x^2 + 6x$      |
| <b>e</b> $x^2 - 2x + 7$ | <b>f</b> $x^2 + 3x - 2$  |
- 2 Write the following quadratic expressions in the form  $p(x + q)^2 + r$
- |                           |                           |
|---------------------------|---------------------------|
| <b>a</b> $2x^2 - 8x - 16$ | <b>b</b> $4x^2 - 8x - 16$ |
| <b>c</b> $3x^2 + 12x - 9$ | <b>d</b> $2x^2 + 6x - 8$  |
- 3 Complete the square.
- |                          |                          |
|--------------------------|--------------------------|
| <b>a</b> $2x^2 + 3x + 6$ | <b>b</b> $3x^2 - 2x$     |
| <b>c</b> $5x^2 + 3x$     | <b>d</b> $3x^2 + 5x + 3$ |

## Extend

- 4 Write  $(25x^2 + 30x + 12)$  in the form  $(ax + b)^2 + c$ .

## Answers

- 1
- |                           |  |
|---------------------------|--|
| <b>a</b> $(x + 2)^2 - 1$  | <b>b</b> $(x - 5)^2 - 28$                                |
| <b>c</b> $(x - 4)^2 - 16$ | <b>d</b> $(x + 3)^2 - 9$                                 |
| <b>e</b> $(x - 1)^2 + 6$  | <b>f</b> $\left(x + \frac{3}{2}\right)^2 - \frac{17}{4}$ |
- 2
- |                            |   |
|----------------------------|---|
| <b>a</b> $2(x - 2)^2 - 24$ | <b>b</b> $4(x - 1)^2 - 20$                                |
| <b>c</b> $3(x + 2)^2 - 21$ | <b>d</b> $2\left(x + \frac{3}{2}\right)^2 - \frac{25}{2}$ |
- 3
- |  |  |
|--|--|
| <b>a</b> $2\left(x + \frac{3}{4}\right)^2 + \frac{39}{8}$  | <b>b</b> $3\left(x - \frac{1}{3}\right)^2 - \frac{1}{3}$   |
| <b>c</b> $5\left(x + \frac{3}{10}\right)^2 - \frac{9}{20}$ | <b>d</b> $3\left(x + \frac{5}{6}\right)^2 + \frac{11}{12}$ |
- 4  $(5x + 3)^2 + 3$

# Solving quadratics by factorisation

## Key points

- A quadratic equation is an equation in the form  $ax^2 + bx + c = 0$  where  $a \neq 0$ .
- To factorise a quadratic equation find two numbers whose sum is  $b$  and whose products is  $ac$ .
- When the product of two numbers is 0, then at least one of the numbers must be 0.
- If a quadratic can be solved it will have two solutions (these may be equal).

## Examples

**Example 1** Solve  $5x^2 = 15x$

$5x^2 = 15x$ $5x^2 - 15x = 0$ $5x(x - 3) = 0$ <p>So <math>5x = 0</math> or <math>(x - 3) = 0</math></p> <p>Therefore <math>x = 0</math> or <math>x = 3</math></p>	<ol style="list-style-type: none"> <li>1 Rearrange the equation so that all of the terms are on one side of the equation and it is equal to zero. Do not divide both sides by <math>x</math> as this would lose the solution <math>x = 0</math>.</li> <li>2 Factorise the quadratic equation. <math>5x</math> is a common factor.</li> <li>3 When two values multiply to make zero, at least one of the values must be zero.</li> <li>4 Solve these two equations.</li> </ol>
---	---

**Example 2** Solve  $x^2 + 7x + 12 = 0$

$x^2 + 7x + 12 = 0$ $b = 7, ac = 12$ $x^2 + 4x + 3x + 12 = 0$ $x(x + 4) + 3(x + 4) = 0$ $(x + 4)(x + 3) = 0$ <p>So <math>(x + 4) = 0</math> or <math>(x + 3) = 0</math></p> <p>Therefore <math>x = -4</math> or <math>x = -3</math></p>	<ol style="list-style-type: none"> <li>1 Factorise the quadratic equation. Work out the two factors of <math>ac = 12</math> which add to give you <math>b = 7</math>. (4 and 3)</li> <li>2 Rewrite the <math>b</math> term (<math>7x</math>) using these two factors.</li> <li>3 Factorise the first two terms and the last two terms.</li> <li>4 <math>(x + 4)</math> is a factor of both terms.</li> <li>5 When two values multiply to make zero, at least one of the values must be zero.</li> <li>6 Solve these two equations.</li> </ol>
---	---

**Example 3** Solve  $9x^2 - 16 = 0$

$9x^2 - 16 = 0$ $(3x + 4)(3x - 4) = 0$ <p>So <math>(3x + 4) = 0</math> or <math>(3x - 4) = 0</math></p> $x = -\frac{4}{3} \text{ or } x = \frac{4}{3}$	<ol style="list-style-type: none"> <li>1 Factorise the quadratic equation. This is the difference of two squares as the two terms are <math>(3x)^2</math> and <math>(4)^2</math>.</li> <li>2 When two values multiply to make zero, at least one of the values must be zero.</li> <li>3 Solve these two equations.</li> </ol>
--	---

**Example 4** Solve  $2x^2 - 5x - 12 = 0$

$b = -5, ac = -24$ <p>So <math>2x^2 - 8x + 3x - 12 = 0</math></p> $2x(x - 4) + 3(x - 4) = 0$ $(x - 4)(2x + 3) = 0$ <p>So <math>(x - 4) = 0</math> or <math>(2x + 3) = 0</math></p> $x = 4 \text{ or } x = -\frac{3}{2}$	<ol style="list-style-type: none"> <li><b>1</b> Factorise the quadratic equation. Work out the two factors of <math>ac = -24</math> which add to give you <math>b = -5</math>. (-8 and 3)</li> <li><b>2</b> Rewrite the <math>b</math> term (<math>-5x</math>) using these two factors.</li> <li><b>3</b> Factorise the first two terms and the last two terms.</li> <li><b>4</b> <math>(x - 4)</math> is a factor of both terms.</li> <li><b>5</b> When two values multiply to make zero, at least one of the values must be zero.</li> <li><b>6</b> Solve these two equations.</li> </ol>
---	---

## Practice

**1** Solve

- |   |  |
|---|--|
| <p><b>a</b> <math>6x^2 + 4x = 0</math></p> <p><b>c</b> <math>x^2 + 7x + 10 = 0</math></p> <p><b>e</b> <math>x^2 - 3x - 4 = 0</math></p> <p><b>g</b> <math>x^2 - 10x + 24 = 0</math></p> <p><b>i</b> <math>x^2 + 3x - 28 = 0</math></p> <p><b>k</b> <math>2x^2 - 7x - 4 = 0</math></p> | <p><b>b</b> <math>28x^2 - 21x = 0</math></p> <p><b>d</b> <math>x^2 - 5x + 6 = 0</math></p> <p><b>f</b> <math>x^2 + 3x - 10 = 0</math></p> <p><b>h</b> <math>x^2 - 36 = 0</math></p> <p><b>j</b> <math>x^2 - 6x + 9 = 0</math></p> <p><b>l</b> <math>3x^2 - 13x - 10 = 0</math></p> |
|---|--|

**2** Solve

- |   |  |
|---|--|
| <p><b>a</b> <math>x^2 - 3x = 10</math></p> <p><b>c</b> <math>x^2 + 5x = 24</math></p> <p><b>e</b> <math>x(x + 2) = 2x + 25</math></p> <p><b>g</b> <math>x(3x + 1) = x^2 + 15</math></p> | <p><b>b</b> <math>x^2 - 3 = 2x</math></p> <p><b>d</b> <math>x^2 - 42 = x</math></p> <p><b>f</b> <math>x^2 - 30 = 3x - 2</math></p> <p><b>h</b> <math>3x(x - 1) = 2(x + 1)</math></p> |
|---|--|

**Hint**

Get all terms onto one side of the equation.

## Answers

- |   |  |
|---|--|
| <p><b>1 a</b> <math>x = 0</math> or <math>x = -\frac{2}{3}</math></p> <p><b>c</b> <math>x = -5</math> or <math>x = -2</math></p> <p><b>e</b> <math>x = -1</math> or <math>x = 4</math></p> <p><b>g</b> <math>x = 4</math> or <math>x = 6</math></p> <p><b>i</b> <math>x = -7</math> or <math>x = 4</math></p> <p><b>k</b> <math>x = -\frac{1}{2}</math> or <math>x = 4</math></p> | <p><b>b</b> <math>x = 0</math> or <math>x = \frac{3}{4}</math></p> <p><b>d</b> <math>x = 2</math> or <math>x = 3</math></p> <p><b>f</b> <math>x = -5</math> or <math>x = 2</math></p> <p><b>h</b> <math>x = -6</math> or <math>x = 6</math></p> <p><b>j</b> <math>x = 3</math></p> <p><b>l</b> <math>x = -\frac{2}{3}</math> or <math>x = 5</math></p> |
| <p><b>2 a</b> <math>x = -2</math> or <math>x = 5</math></p> <p><b>c</b> <math>x = -8</math> or <math>x = 3</math></p> <p><b>e</b> <math>x = -5</math> or <math>x = 5</math></p> <p><b>g</b> <math>x = -3</math> or <math>x = 2\frac{1}{2}</math></p>  | <p><b>b</b> <math>x = -1</math> or <math>x = 3</math></p> <p><b>d</b> <math>x = -6</math> or <math>x = 7</math></p> <p><b>f</b> <math>x = -4</math> or <math>x = 7</math></p> <p><b>h</b> <math>x = -\frac{1}{3}</math> or <math>x = 2</math></p>  |

# Solving quadratics by completing the square

## Key points

- Completing the square lets you write a quadratic equation in the form  $p(x + q)^2 + r = 0$ .

## Examples

**Example 5** Solve  $x^2 + 6x + 4 = 0$ . Give your solutions in surd form.

$x^2 + 6x + 4 = 0$ $(x + 3)^2 - 9 + 4 = 0$ $(x + 3)^2 - 5 = 0$ $(x + 3)^2 = 5$ $x + 3 = \pm\sqrt{5}$ $x = \pm\sqrt{5} - 3$ <p>So <math>x = -\sqrt{5} - 3</math> or <math>x = \sqrt{5} - 3</math></p>	<ol style="list-style-type: none"> <li>1 Write <math>x^2 + bx + c = 0</math> in the form <math>\left(x + \frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2 + c = 0</math></li> <li>2 Simplify.</li> <li>3 Rearrange the equation to work out <math>x</math>. First, add 5 to both sides.</li> <li>4 Square root both sides. Remember that the square root of a value gives two answers.</li> <li>5 Subtract 3 from both sides to solve the equation.</li> <li>6 Write down both solutions.</li> </ol>
--	---

**Example 6** Solve  $2x^2 - 7x + 4 = 0$ . Give your solutions in surd form.

$2x^2 - 7x + 4 = 0$ $2\left(x^2 - \frac{7}{2}x\right) + 4 = 0$ $2\left[\left(x - \frac{7}{4}\right)^2 - \left(\frac{7}{4}\right)^2\right] + 4 = 0$ $2\left(x - \frac{7}{4}\right)^2 - \frac{49}{8} + 4 = 0$ $2\left(x - \frac{7}{4}\right)^2 - \frac{17}{8} = 0$ $2\left(x - \frac{7}{4}\right)^2 = \frac{17}{8}$	<ol style="list-style-type: none"> <li>1 Before completing the square write <math>ax^2 + bx + c</math> in the form <math>a\left(x^2 + \frac{b}{a}x\right) + c</math></li> <li>2 Now complete the square by writing <math>x^2 - \frac{7}{2}x</math> in the form <math>\left(x + \frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2</math></li> <li>3 Expand the square brackets.</li> <li>4 Simplify.</li> </ol> <p style="text-align: right;"><i>(continued on next page)</i></p> <ol style="list-style-type: none"> <li>5 Rearrange the equation to work out <math>x</math>. First, add <math>\frac{17}{8}</math> to both sides.</li> </ol>
---	---

$\left(x - \frac{7}{4}\right)^2 = \frac{17}{16}$ $x - \frac{7}{4} = \pm \frac{\sqrt{17}}{4}$ $x = \pm \frac{\sqrt{17}}{4} + \frac{7}{4}$ <p>So <math>x = \frac{7}{4} - \frac{\sqrt{17}}{4}</math> or <math>x = \frac{7}{4} + \frac{\sqrt{17}}{4}</math></p>	<p><b>6</b> Divide both sides by 2.</p> <p><b>7</b> Square root both sides. Remember that the square root of a value gives two answers.</p> <p><b>8</b> Add <math>\frac{7}{4}</math> to both sides.</p> <p><b>9</b> Write down both the solutions.</p>
---	--

## Practice

**3** Solve by completing the square.

**a**  $x^2 - 4x - 3 = 0$

**c**  $x^2 + 8x - 5 = 0$

**e**  $2x^2 + 8x - 5 = 0$

**b**  $x^2 - 10x + 4 = 0$

**d**  $x^2 - 2x - 6 = 0$

**f**  $5x^2 + 3x - 4 = 0$

**4** Solve by completing the square.

**a**  $(x - 4)(x + 2) = 5$

**b**  $2x^2 + 6x - 7 = 0$

**c**  $x^2 - 5x + 3 = 0$

### Hint

Get all terms onto one side of the equation.

## Answers

**3 a**  $x = 2 + \sqrt{7}$  or  $x = 2 - \sqrt{7}$

**c**  $x = -4 + \sqrt{21}$  or  $x = -4 - \sqrt{21}$

**e**  $x = -2 + \sqrt{6.5}$  or  $x = -2 - \sqrt{6.5}$

**b**  $x = 5 + \sqrt{21}$  or  $x = 5 - \sqrt{21}$

**d**  $x = 1 + \sqrt{7}$  or  $x = 1 - \sqrt{7}$

**f**  $x = \frac{-3 + \sqrt{89}}{10}$  or  $x = \frac{-3 - \sqrt{89}}{10}$

**4 a**  $x = 1 + \sqrt{14}$  or  $x = 1 - \sqrt{14}$

**c**  $x = \frac{5 + \sqrt{13}}{2}$  or  $x = \frac{5 - \sqrt{13}}{2}$

**b**  $x = \frac{-3 + \sqrt{23}}{2}$  or  $x = \frac{-3 - \sqrt{23}}{2}$

# Solving quadratics by using the formula

## Key points

- Any quadratic equation of the form  $ax^2 + bx + c = 0$  can be solved using the formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

- If  $b^2 - 4ac$  is negative then the quadratic equation does not have any real solutions.
- It is useful to write down the formula before substituting the values for  $a$ ,  $b$  and  $c$ .

## Examples

**Example 7** Solve  $x^2 + 6x + 4 = 0$ . Give your solutions in surd form.

$a = 1, b = 6, c = 4$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-6 \pm \sqrt{6^2 - 4(1)(4)}}{2(1)}$ $x = \frac{-6 \pm \sqrt{20}}{2}$ $x = \frac{-6 \pm 2\sqrt{5}}{2}$ $x = -3 \pm \sqrt{5}$ <p>So <math>x = -3 - \sqrt{5}</math> or <math>x = \sqrt{5} - 3</math></p>	<ol style="list-style-type: none"> <li>Identify <math>a</math>, <math>b</math> and <math>c</math> and write down the formula. Remember that <math>-b \pm \sqrt{b^2 - 4ac}</math> is all over <math>2a</math>, not just part of it.</li> <li>Substitute <math>a = 1</math>, <math>b = 6</math>, <math>c = 4</math> into the formula.</li> <li>Simplify. The denominator is 2, but this is only because <math>a = 1</math>. The denominator will not always be 2.</li> <li>Simplify <math>\sqrt{20}</math>. <math>\sqrt{20} = \sqrt{4 \times 5} = \sqrt{4} \times \sqrt{5} = 2\sqrt{5}</math></li> <li>Simplify by dividing numerator and denominator by 2.</li> <li>Write down both the solutions.</li> </ol>
---	--

**Example 8** Solve  $3x^2 - 7x - 2 = 0$ . Give your solutions in surd form.

$a = 3, b = -7, c = -2$ $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-(-7) \pm \sqrt{(-7)^2 - 4(3)(-2)}}{2(3)}$ $x = \frac{7 \pm \sqrt{73}}{6}$ <p>So <math>x = \frac{7 - \sqrt{73}}{6}</math> or <math>x = \frac{7 + \sqrt{73}}{6}</math></p>	<ol style="list-style-type: none"> <li>Identify <math>a</math>, <math>b</math> and <math>c</math>, making sure you get the signs right and write down the formula. Remember that <math>-b \pm \sqrt{b^2 - 4ac}</math> is all over <math>2a</math>, not just part of it.</li> <li>Substitute <math>a = 3</math>, <math>b = -7</math>, <math>c = -2</math> into the formula.</li> <li>Simplify. The denominator is 6 when <math>a = 3</math>. A common mistake is to always write a denominator of 2.</li> <li>Write down both the solutions.</li> </ol>
---	--

## Practice

5 Solve, giving your solutions in surd form.

**a**  $3x^2 + 6x + 2 = 0$

**b**  $2x^2 - 4x - 7 = 0$

6 Solve the equation  $x^2 - 7x + 2 = 0$

Give your solutions in the form  $\frac{a \pm \sqrt{b}}{c}$ , where  $a$ ,  $b$  and  $c$  are integers.

7 Solve  $10x^2 + 3x + 3 = 5$

Give your solution in surd form.

### Hint

Get all terms onto one side of the equation.

## Extend

8 Choose an appropriate method to solve each quadratic equation, giving your answer in surd form when necessary.

**a**  $4x(x - 1) = 3x - 2$

**b**  $10 = (x + 1)^2$

**c**  $x(3x - 1) = 10$

## Answers

5 **a**  $x = -1 + \frac{\sqrt{3}}{3}$  or  $x = -1 - \frac{\sqrt{3}}{3}$       **b**  $x = 1 + \frac{3\sqrt{2}}{2}$  or  $x = 1 - \frac{3\sqrt{2}}{2}$

6  $x = \frac{7 + \sqrt{41}}{2}$  or  $x = \frac{7 - \sqrt{41}}{2}$

7  $x = \frac{-3 + \sqrt{89}}{20}$  or  $x = \frac{-3 - \sqrt{89}}{20}$

8 **a**  $x = \frac{7 + \sqrt{17}}{8}$  or  $x = \frac{7 - \sqrt{17}}{8}$

**b**  $x = -1 + \sqrt{10}$  or  $x = -1 - \sqrt{10}$

**c**  $x = -1\frac{2}{3}$  or  $x = 2$



# Solving linear simultaneous equations using the elimination method

## Key points

- Two equations are simultaneous when they are both true at the same time.
- Solving simultaneous linear equations in two unknowns involves finding the value of each unknown which works for both equations.
- Make sure that the coefficient of one of the unknowns is the same in both equations.
- Eliminate this equal unknown by either subtracting or adding the two equations.

## Examples

**Example 1** Solve the simultaneous equations  $3x + y = 5$  and  $x + y = 1$

$\begin{array}{r} 3x + y = 5 \\ - \quad x + y = 1 \\ \hline 2x \quad = 4 \\ \text{So } x = 2 \end{array}$ <p>Using <math>x + y = 1</math></p> $\begin{array}{r} 2 + y = 1 \\ \text{So } y = -1 \end{array}$ <p>Check:</p> <p>equation 1: <math>3 \times 2 + (-1) = 5</math> YES  equation 2: <math>2 + (-1) = 1</math> YES</p>	<ol style="list-style-type: none"> <li><b>1</b> Subtract the second equation from the first equation to eliminate the <math>y</math> term.</li> <li><b>2</b> To find the value of <math>y</math>, substitute <math>x = 2</math> into one of the original equations.</li> <li><b>3</b> Substitute the values of <math>x</math> and <math>y</math> into both equations to check your answers.</li> </ol>
--	--

**Example 2** Solve  $x + 2y = 13$  and  $5x - 2y = 5$  simultaneously.

$\begin{array}{r} x + 2y = 13 \\ + \quad 5x - 2y = 5 \\ \hline 6x \quad = 18 \\ \text{So } x = 3 \end{array}$ <p>Using <math>x + 2y = 13</math></p> $\begin{array}{r} 3 + 2y = 13 \\ \text{So } y = 5 \end{array}$ <p>Check:</p> <p>equation 1: <math>3 + 2 \times 5 = 13</math> YES  equation 2: <math>5 \times 3 - 2 \times 5 = 5</math> YES</p>	<ol style="list-style-type: none"> <li><b>1</b> Add the two equations together to eliminate the <math>y</math> term.</li> <li><b>2</b> To find the value of <math>y</math>, substitute <math>x = 3</math> into one of the original equations.</li> <li><b>3</b> Substitute the values of <math>x</math> and <math>y</math> into both equations to check your answers.</li> </ol>
--	--

**Example 3** Solve  $2x + 3y = 2$  and  $5x + 4y = 12$  simultaneously.

$\begin{array}{r} (2x + 3y = 2) \times 4 \rightarrow 8x + 12y = 8 \\ (5x + 4y = 12) \times 3 \rightarrow \frac{15x + 12y = 36}{7x = 28} \end{array}$ <p>So <math>x = 4</math></p> <p>Using <math>2x + 3y = 2</math>  <math>2 \times 4 + 3y = 2</math>          So <math>y = -2</math></p> <p>Check:          equation 1: <math>2 \times 4 + 3 \times (-2) = 2</math> YES          equation 2: <math>5 \times 4 + 4 \times (-2) = 12</math> YES</p>	<p><b>1</b> Multiply the first equation by 4 and the second equation by 3 to make the coefficient of <math>y</math> the same for both equations. Then subtract the first equation from the second equation to eliminate the <math>y</math> term.</p> <p><b>2</b> To find the value of <math>y</math>, substitute <math>x = 4</math> into one of the original equations.</p> <p><b>3</b> Substitute the values of <math>x</math> and <math>y</math> into both equations to check your answers.</p>
--	---

## Practice

Solve these simultaneous equations.

**1**  $4x + y = 8$   
 $x + y = 5$

**2**  $3x + y = 7$   
 $3x + 2y = 5$

**3**  $4x + y = 3$   
 $3x - y = 11$

**4**  $3x + 4y = 7$   
 $x - 4y = 5$

**5**  $2x + y = 11$   
 $x - 3y = 9$

**6**  $2x + 3y = 11$   
 $3x + 2y = 4$

## Answers

**1**  $x = 1, y = 4$

**2**  $x = 3, y = -2$

**3**  $x = 2, y = -5$

**4**  $x = 3, y = -\frac{1}{2}$

**5**  $x = 6, y = -1$

**6**  $x = -2, y = 5$

# Solving linear simultaneous equations using the substitution method

## Key points

- The substitution method is the method most commonly used for A level. This is because it is the method used to solve linear and quadratic simultaneous equations.

## Examples

**Example 4** Solve the simultaneous equations  $y = 2x + 1$  and  $5x + 3y = 14$

$5x + 3(2x + 1) = 14$ $5x + 6x + 3 = 14$ $11x + 3 = 14$ $11x = 11$ <p>So <math>x = 1</math></p> <p>Using <math>y = 2x + 1</math></p> $y = 2 \times 1 + 1$ <p>So <math>y = 3</math></p> <p>Check:</p> <p>equation 1: <math>3 = 2 \times 1 + 1</math>      YES</p> <p>equation 2: <math>5 \times 1 + 3 \times 3 = 14</math>    YES</p>	<ol style="list-style-type: none"> <li>Substitute <math>2x + 1</math> for <math>y</math> into the second equation.</li> <li>Expand the brackets and simplify.</li> <li>Work out the value of <math>x</math>.</li> <li>To find the value of <math>y</math>, substitute <math>x = 1</math> into one of the original equations.</li> <li>Substitute the values of <math>x</math> and <math>y</math> into both equations to check your answers.</li> </ol>
--	--

**Example 5** Solve  $2x - y = 16$  and  $4x + 3y = -3$  simultaneously.

$y = 2x - 16$ $4x + 3(2x - 16) = -3$ $4x + 6x - 48 = -3$ $10x - 48 = -3$ $10x = 45$ <p>So <math>x = 4\frac{1}{2}</math></p> <p>Using <math>y = 2x - 16</math></p> $y = 2 \times 4\frac{1}{2} - 16$ <p>So <math>y = -7</math></p> <p>Check:</p> <p>equation 1: <math>2 \times 4\frac{1}{2} - (-7) = 16</math>      YES</p> <p>equation 2: <math>4 \times 4\frac{1}{2} + 3 \times (-7) = -3</math>    YES</p>	<ol style="list-style-type: none"> <li>Rearrange the first equation.</li> <li>Substitute <math>2x - 16</math> for <math>y</math> into the second equation.</li> <li>Expand the brackets and simplify.</li> <li>Work out the value of <math>x</math>.</li> <li>To find the value of <math>y</math>, substitute <math>x = 4\frac{1}{2}</math> into one of the original equations.</li> <li>Substitute the values of <math>x</math> and <math>y</math> into both equations to check your answers.</li> </ol>
---	---

## Practice

Solve these simultaneous equations.

7  $y = x - 4$   
 $2x + 5y = 43$

8  $y = 2x - 3$   
 $5x - 3y = 11$

9  $2y = 4x + 5$   
 $9x + 5y = 22$

10  $2x = y - 2$   
 $8x - 5y = -11$

11  $3x + 4y = 8$   
 $2x - y = -13$

12  $3y = 4x - 7$   
 $2y = 3x - 4$

13  $3x = y - 1$   
 $2y - 2x = 3$

14  $3x + 2y + 1 = 0$   
 $4y = 8 - x$

## Extend

15 Solve the simultaneous equations  $3x + 5y - 20 = 0$  and  $2(x + y) = \frac{3(y - x)}{4}$ .

## Answers

7  $x = 9, y = 5$

8  $x = -2, y = -7$

9  $x = \frac{1}{2}, y = 3\frac{1}{2}$

10  $x = \frac{1}{2}, y = 3$

11  $x = -4, y = 5$

12  $x = -2, y = -5$

13  $x = \frac{1}{4}, y = 1\frac{3}{4}$

14  $x = -2, y = 2\frac{1}{2}$

15  $x = -2\frac{1}{2}, y = 5\frac{1}{2}$

# Solving linear and quadratic simultaneous equations

## Key points

- Make one of the unknowns the subject of the linear equation (rearranging where necessary).
- Use the linear equation to substitute into the quadratic equation.
- There are usually two pairs of solutions.

## Examples

**Example 1** Solve the simultaneous equations  $y = x + 1$  and  $x^2 + y^2 = 13$

$x^2 + (x + 1)^2 = 13$ $x^2 + x^2 + x + x + 1 = 13$ $2x^2 + 2x + 1 = 13$ $2x^2 + 2x - 12 = 0$ $(2x - 4)(x + 3) = 0$ <p>So <math>x = 2</math> or <math>x = -3</math></p> <p>Using <math>y = x + 1</math>            When <math>x = 2</math>, <math>y = 2 + 1 = 3</math>            When <math>x = -3</math>, <math>y = -3 + 1 = -2</math></p> <p>So the solutions are  <math>x = 2, y = 3</math> and <math>x = -3, y = -2</math></p> <p>Check:</p> <p>equation 1: <math>3 = 2 + 1</math> YES            and <math>-2 = -3 + 1</math> YES</p> <p>equation 2: <math>2^2 + 3^2 = 13</math> YES            and <math>(-3)^2 + (-2)^2 = 13</math> YES</p>	<ol style="list-style-type: none"> <li>1 Substitute <math>x + 1</math> for <math>y</math> into the second equation.</li> <li>2 Expand the brackets and simplify.</li> <li>3 Factorise the quadratic equation.</li> <li>4 Work out the values of <math>x</math>.</li> <li>5 To find the value of <math>y</math>, substitute both values of <math>x</math> into one of the original equations.</li> <li>6 Substitute both pairs of values of <math>x</math> and <math>y</math> into both equations to check your answers.</li> </ol>
--	--

**Example 2** Solve  $2x + 3y = 5$  and  $2y^2 + xy = 12$  simultaneously.

$x = \frac{5-3y}{2}$ $2y^2 + \left(\frac{5-3y}{2}\right)y = 12$ $2y^2 + \frac{5y-3y^2}{2} = 12$ $4y^2 + 5y - 3y^2 = 24$ $y^2 + 5y - 24 = 0$ $(y+8)(y-3) = 0$ <p>So <math>y = -8</math> or <math>y = 3</math></p> <p>Using <math>2x + 3y = 5</math>              When <math>y = -8</math>, <math>2x + 3 \times (-8) = 5</math>, <math>x = 14.5</math>              When <math>y = 3</math>, <math>2x + 3 \times 3 = 5</math>, <math>x = -2</math></p> <p>So the solutions are  <math>x = 14.5</math>, <math>y = -8</math> and <math>x = -2</math>, <math>y = 3</math></p> <p>Check:              equation 1: <math>2 \times 14.5 + 3 \times (-8) = 5</math> YES                                and <math>2 \times (-2) + 3 \times 3 = 5</math> YES              equation 2: <math>2 \times (-8)^2 + 14.5 \times (-8) = 12</math> YES                                and <math>2 \times (3)^2 + (-2) \times 3 = 12</math> YES</p>	<ol style="list-style-type: none"> <li>1 Rearrange the first equation.</li> <li>2 Substitute <math>\frac{5-3y}{2}</math> for <math>x</math> into the second equation. Notice how it is easier to substitute for <math>x</math> than for <math>y</math>.</li> <li>3 Expand the brackets and simplify.</li> <li>4 Factorise the quadratic equation.</li> <li>5 Work out the values of <math>y</math>.</li> <li>6 To find the value of <math>x</math>, substitute both values of <math>y</math> into one of the original equations.</li> <li>7 Substitute both pairs of values of <math>x</math> and <math>y</math> into both equations to check your answers.</li> </ol>
---	--

## Practice

Solve these simultaneous equations.

- |   |   |
|---|---|
| <p><b>1</b> <math>y = 2x + 1</math><br/><math>x^2 + y^2 = 10</math></p>   | <p><b>2</b> <math>y = 6 - x</math><br/><math>x^2 + y^2 = 20</math></p>    |
| <p><b>3</b> <math>y = x - 3</math><br/><math>x^2 + y^2 = 5</math></p>     | <p><b>4</b> <math>y = 9 - 2x</math><br/><math>x^2 + y^2 = 17</math></p>   |
| <p><b>5</b> <math>y = 3x - 5</math><br/><math>y = x^2 - 2x + 1</math></p> | <p><b>6</b> <math>y = x - 5</math><br/><math>y = x^2 - 5x - 12</math></p> |
| <p><b>7</b> <math>y = x + 5</math><br/><math>x^2 + y^2 = 25</math></p>    | <p><b>8</b> <math>y = 2x - 1</math><br/><math>x^2 + xy = 24</math></p>    |
| <p><b>9</b> <math>y = 2x</math><br/><math>y^2 - xy = 8</math></p>         | <p><b>10</b> <math>2x + y = 11</math><br/><math>xy = 15</math></p>        |

## Extend

- |  |   |
|--|---|
| <p><b>11</b> <math>x - y = 1</math><br/><math>x^2 + y^2 = 3</math></p> | <p><b>12</b> <math>y - x = 2</math><br/><math>x^2 + xy = 3</math></p> |
|--|---|

## Answers

**1**  $x = 1, y = 3$

$$x = -\frac{9}{5}, y = -\frac{13}{5}$$

**2**  $x = 2, y = 4$

$$x = 4, y = 2$$

**3**  $x = 1, y = -2$

$$x = 2, y = -1$$

**4**  $x = 4, y = 1$

$$x = \frac{16}{5}, y = \frac{13}{5}$$

**5**  $x = 3, y = 4$

$$x = 2, y = 1$$

**6**  $x = 7, y = 2$

$$x = -1, y = -6$$

**7**  $x = 0, y = 5$

$$x = -5, y = 0$$

**8**  $x = -\frac{8}{3}, y = -\frac{19}{3}$

$$x = 3, y = 5$$

**9**  $x = -2, y = -4$

$$x = 2, y = 4$$

**10**  $x = \frac{5}{2}, y = 6$

$$x = 3, y = 5$$

**11**  $x = \frac{1+\sqrt{5}}{2}, y = \frac{-1+\sqrt{5}}{2}$

$$x = \frac{1-\sqrt{5}}{2}, y = \frac{-1-\sqrt{5}}{2}$$

**12**  $x = \frac{-1+\sqrt{7}}{2}, y = \frac{3+\sqrt{7}}{2}$


$$x = \frac{-1-\sqrt{7}}{2}, y = \frac{3-\sqrt{7}}{2}$$

# Straight line graphs

## Key points

- A straight line has the equation  $y = mx + c$ , where  $m$  is the gradient and  $c$  is the  $y$ -intercept (where  $x = 0$ ).
- The equation of a straight line can be written in the form  $ax + by + c = 0$ , where  $a$ ,  $b$  and  $c$  are integers.
- When given the coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$  of two points on a line the gradient is calculated using the

$$\text{formula } m = \frac{y_2 - y_1}{x_2 - x_1}$$



## Examples

**Example 1** A straight line has gradient  $-\frac{1}{2}$  and  $y$ -intercept 3.

Write the equation of the line in the form  $ax + by + c = 0$ .

$m = -\frac{1}{2} \text{ and } c = 3$ $\text{So } y = -\frac{1}{2}x + 3$ $\frac{1}{2}x + y - 3 = 0$ $x + 2y - 6 = 0$	<ol style="list-style-type: none"> <li><b>1</b> A straight line has equation <math>y = mx + c</math>. Substitute the gradient and <math>y</math>-intercept given in the question into this equation.</li> <li><b>2</b> Rearrange the equation so all the terms are on one side and 0 is on the other side.</li> <li><b>3</b> Multiply both sides by 2 to eliminate the denominator.</li> </ol>
--	--

**Example 2** Find the gradient and the  $y$ -intercept of the line with the equation  $3y - 2x + 4 = 0$ .

$3y - 2x + 4 = 0$ $3y = 2x - 4$ $y = \frac{2}{3}x - \frac{4}{3}$ $\text{Gradient } = m = \frac{2}{3}$ $\text{y-intercept } = c = -\frac{4}{3}$	<ol style="list-style-type: none"> <li><b>1</b> Make <math>y</math> the subject of the equation.</li> <li><b>2</b> Divide all the terms by three to get the equation in the form <math>y = \dots</math></li> <li><b>3</b> In the form <math>y = mx + c</math>, the gradient is <math>m</math> and the <math>y</math>-intercept is <math>c</math>.</li> </ol>
--	--



**Example 3** Find the equation of the line which passes through the point (5, 13) and has gradient 3.

$m = 3$ $y = 3x + c$ $13 = 3 \times 5 + c$ $13 = 15 + c$ $c = -2$ $y = 3x - 2$	<ol style="list-style-type: none"> <li><b>1</b> Substitute the gradient given in the question into the equation of a straight line <math>y = mx + c</math>.</li> <li><b>2</b> Substitute the coordinates <math>x = 5</math> and <math>y = 13</math> into the equation.</li> <li><b>3</b> Simplify and solve the equation.</li> <li><b>4</b> Substitute <math>c = -2</math> into the equation <math>y = 3x + c</math></li> </ol>
--	---

**Example 4** Find the equation of the line passing through the points with coordinates (2, 4) and (8, 7).

$x_1 = 2, x_2 = 8, y_1 = 4 \text{ and } y_2 = 7$ $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - 4}{8 - 2} = \frac{3}{6} = \frac{1}{2}$ $y = \frac{1}{2}x + c$ $4 = \frac{1}{2} \times 2 + c$ $c = 3$ $y = \frac{1}{2}x + 3$	<ol style="list-style-type: none"> <li><b>1</b> Substitute the coordinates into the equation <math>m = \frac{y_2 - y_1}{x_2 - x_1}</math> to work out the gradient of the line.</li> <li><b>2</b> Substitute the gradient into the equation of a straight line <math>y = mx + c</math>.</li> <li><b>3</b> Substitute the coordinates of either point into the equation.</li> <li><b>4</b> Simplify and solve the equation.</li> <li><b>5</b> Substitute <math>c = 3</math> into the equation <math>y = \frac{1}{2}x + c</math></li> </ol>
--	---

## Practice

1 Find the gradient and the y-intercept of the following equations.

- a**  $y = 3x + 5$                       **b**  $y = -\frac{1}{2}x - 7$   
**c**  $2y = 4x - 3$                       **d**  $x + y = 5$   
**e**  $2x - 3y - 7 = 0$                   **f**  $5x + y - 4 = 0$

**Hint**  
Rearrange the equations to the form  $y = mx + c$

2 Copy and complete the table, giving the equation of the line in the form  $y = mx + c$ .

Gradient	y-intercept	Equation of the line
5	0	
-3	2	
4	-7	

3 Find, in the form  $ax + by + c = 0$  where  $a$ ,  $b$  and  $c$  are integers, an equation for each of the lines with the following gradients and y-intercepts.

- a** gradient  $-\frac{1}{2}$ , y-intercept  $-7$               **b** gradient 2, y-intercept 0  
**c** gradient  $\frac{2}{3}$ , y-intercept 4                  **d** gradient  $-1.2$ , y-intercept  $-2$

4 Write an equation for the line which passes through the point (2, 5) and has gradient 4.

5 Write an equation for the line which passes through the point (6, 3) and has gradient  $-\frac{2}{3}$

6 Write an equation for the line passing through each of the following pairs of points.

- a** (4, 5), (10, 17)                              **b** (0, 6), (-4, 8)  
**c** (-1, -7), (5, 23)                           **d** (3, 10), (4, 7)

## Extend

7 The equation of a line is  $2y + 3x - 6 = 0$ .  
Write as much information as possible about this line.

**Answers**

- 1 a**  $m = 3, c = 5$                       **b**  $m = -\frac{1}{2}, c = -7$   
**c**  $m = 2, c = -\frac{3}{2}$                       **d**  $m = -1, c = 5$   
**e**  $m = \frac{2}{3}, c = -\frac{7}{3}$  or  $-2\frac{1}{3}$                       **f**  $m = -5, c = 4$

**2**

Gradient	y-intercept	Equation of the line
5	0	$y = 5x$
-3	2	$y = -3x + 2$
4	-7	$y = 4x - 7$

- 3 a**  $x + 2y + 14 = 0$                       **b**  $2x - y = 0$   
**c**  $2x - 3y + 12 = 0$                       **d**  $6x + 5y + 10 = 0$

**4**  $y = 4x - 3$

**5**  $y = -\frac{2}{3}x + 7$

**6 a**  $y = 2x - 3$                       **b**  $y = -\frac{1}{2}x + 6$

**c**  $y = 5x - 2$                       **d**  $y = -3x + 19$

**7**  $y = -\frac{3}{2}x + 3$ , the gradient is  $-\frac{3}{2}$  and the y-intercept is 3.

The line intercepts the axes at (0, 3) and (2, 0).

Students may sketch the line or give coordinates that lie on the line such as  $\left(1, \frac{3}{2}\right)$  or (4, -3).


# Parallel and perpendicular lines

## A LEVEL LINKS

Scheme of work: 2a. Straight-line graphs, parallel/perpendicular, length and area problems

### Key points

- When lines are parallel they have the same gradient.
- A line perpendicular to the line with equation  $y = mx + c$  has gradient  $-\frac{1}{m}$ .



### Examples

**Example 1** Find the equation of the line parallel to  $y = 2x + 4$  which passes through the point  $(4, 9)$ .

$y = 2x + 4$ $m = 2$ $y = 2x + c$ $9 = 2 \times 4 + c$ $9 = 8 + c$ $c = 1$ $y = 2x + 1$	<ol style="list-style-type: none"> <li>1 As the lines are parallel they have the same gradient.</li> <li>2 Substitute <math>m = 2</math> into the equation of a straight line <math>y = mx + c</math>.</li> <li>3 Substitute the coordinates into the equation <math>y = 2x + c</math></li> <li>4 Simplify and solve the equation.</li> <li>5 Substitute <math>c = 1</math> into the equation <math>y = 2x + c</math></li> </ol>
---	--

**Example 2** Find the equation of the line perpendicular to  $y = 2x - 3$  which passes through the point  $(-2, 5)$ .

$y = 2x - 3$ $m = 2$ $-\frac{1}{m} = -\frac{1}{2}$ $y = -\frac{1}{2}x + c$ $5 = -\frac{1}{2} \times (-2) + c$ $5 = 1 + c$ $c = 4$ $y = -\frac{1}{2}x + 4$	<ol style="list-style-type: none"> <li>1 As the lines are perpendicular, the gradient of the perpendicular line is <math>-\frac{1}{m}</math>.</li> <li>2 Substitute <math>m = -\frac{1}{2}</math> into <math>y = mx + c</math>.</li> <li>3 Substitute the coordinates <math>(-2, 5)</math> into the equation <math>y = -\frac{1}{2}x + c</math></li> <li>4 Simplify and solve the equation.</li> <li>5 Substitute <math>c = 4</math> into <math>y = -\frac{1}{2}x + c</math>.</li> </ol>
---	--

**Example 3** A line passes through the points (0, 5) and (9, -1).  
Find the equation of the line which is perpendicular to the line and passes through its midpoint.

$x_1 = 0, x_2 = 9, y_1 = 5 \text{ and } y_2 = -1$ $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-1 - 5}{9 - 0}$ $= \frac{-6}{9} = -\frac{2}{3}$ $-\frac{1}{m} = \frac{3}{2}$ $y = \frac{3}{2}x + c$ $\text{Midpoint} = \left( \frac{0+9}{2}, \frac{5+(-1)}{2} \right) = \left( \frac{9}{2}, 2 \right)$ $2 = \frac{3}{2} \times \frac{9}{2} + c$ $c = -\frac{19}{4}$ $y = \frac{3}{2}x - \frac{19}{4}$	<ol style="list-style-type: none"> <li><b>1</b> Substitute the coordinates into the equation <math>m = \frac{y_2 - y_1}{x_2 - x_1}</math> to work out the gradient of the line.</li> <li><b>2</b> As the lines are perpendicular, the gradient of the perpendicular line is <math>-\frac{1}{m}</math>.</li> <li><b>3</b> Substitute the gradient into the equation <math>y = mx + c</math>.</li> <li><b>4</b> Work out the coordinates of the midpoint of the line.</li> <li><b>5</b> Substitute the coordinates of the midpoint into the equation.</li> <li><b>6</b> Simplify and solve the equation.</li> <li><b>7</b> Substitute <math>c = -\frac{19}{4}</math> into the equation <math>y = \frac{3}{2}x + c</math>.</li> </ol>
--	--

## Practice

- 1** Find the equation of the line parallel to each of the given lines and which passes through each of the given points.
 

<b>a</b> $y = 3x + 1$ (3, 2)	<b>b</b> $y = 3 - 2x$ (1, 3)
<b>c</b> $2x + 4y + 3 = 0$ (6, -3)	<b>d</b> $2y - 3x + 2 = 0$ (8, 20)

- 2** Find the equation of the line perpendicular to  $y = \frac{1}{2}x - 3$  which passes through the point (-5, 3).

**Hint**

If  $m = \frac{a}{b}$  then the negative reciprocal  $-\frac{1}{m} = -\frac{b}{a}$

- 3** Find the equation of the line perpendicular to each of the given lines and which passes through each of the given points.
 

<b>a</b> $y = 2x - 6$ (4, 0)	<b>b</b> $y = -\frac{1}{3}x + \frac{1}{2}$ (2, 13)
<b>c</b> $x - 4y - 4 = 0$ (5, 15)	<b>d</b> $5y + 2x - 5 = 0$ (6, 7)



# Rearranging equations

## Key points

- To change the subject of a formula, get the terms containing the subject on one side and everything else on the other side.
- You may need to factorise the terms containing the new subject.

## Examples

**Example 1** Make  $t$  the subject of the formula  $v = u + at$ .

$v = u + at$ $v - u = at$ $t = \frac{v - u}{a}$	<ol style="list-style-type: none"> <li><b>1</b> Get the terms containing <math>t</math> on one side and everything else on the other side.</li> <li><b>2</b> Divide throughout by <math>a</math>.</li> </ol>
---	--

**Example 2** Make  $t$  the subject of the formula  $r = 2t - \pi t$ .

$r = 2t - \pi t$ $r = t(2 - \pi)$ $t = \frac{r}{2 - \pi}$	<ol style="list-style-type: none"> <li><b>1</b> All the terms containing <math>t</math> are already on one side and everything else is on the other side.</li> <li><b>2</b> Factorise as <math>t</math> is a common factor.</li> <li><b>3</b> Divide throughout by <math>2 - \pi</math>.</li> </ol>
---	---

**Example 3** Make  $t$  the subject of the formula  $\frac{t + r}{5} = \frac{3t}{2}$ .

$\frac{t + r}{5} = \frac{3t}{2}$ $2t + 2r = 15t$ $2r = 13t$ $t = \frac{2r}{13}$	<ol style="list-style-type: none"> <li><b>1</b> Remove the fractions first by multiplying throughout by 10.</li> <li><b>2</b> Get the terms containing <math>t</math> on one side and everything else on the other side and simplify.</li> <li><b>3</b> Divide throughout by 13.</li> </ol>
---	---

**Example 4** Make  $t$  the subject of the formula  $r = \frac{3t+5}{t-1}$ .

$r = \frac{3t+5}{t-1}$ $r(t-1) = 3t+5$ $rt - r = 3t+5$ $rt - 3t = 5+r$ $t(r-3) = 5+r$ $t = \frac{5+r}{r-3}$	<ol style="list-style-type: none"> <li><b>1</b> Remove the fraction first by multiplying throughout by <math>t-1</math>.</li> <li><b>2</b> Expand the brackets.</li> <li><b>3</b> Get the terms containing <math>t</math> on one side and everything else on the other side.</li> <li><b>4</b> Factorise the LHS as <math>t</math> is a common factor.</li> <li><b>5</b> Divide throughout by <math>r-3</math>.</li> </ol>
---	--

## Practice

Change the subject of each formula to the letter given in the brackets.

- |  |  |  |
|--|--|--|
| <b>1</b> $C = \pi d$ [ $d$ ]                       | <b>2</b> $P = 2l + 2w$ [ $w$ ]           | <b>3</b> $D = \frac{S}{T}$ [ $T$ ]       |
| <b>4</b> $p = \frac{q-r}{t}$ [ $t$ ]               | <b>5</b> $u = at - \frac{1}{2}t$ [ $t$ ] | <b>6</b> $V = ax + 4x$ [ $x$ ]           |
| <b>7</b> $\frac{y-7x}{2} = \frac{7-2y}{3}$ [ $y$ ] | <b>8</b> $x = \frac{2a-1}{3-a}$ [ $a$ ]  | <b>9</b> $x = \frac{b-c}{d}$ [ $d$ ]     |
| <b>10</b> $h = \frac{7g-9}{2+g}$ [ $g$ ]           | <b>11</b> $e(9+x) = 2e+1$ [ $e$ ]        | <b>12</b> $y = \frac{2x+3}{4-x}$ [ $x$ ] |

**13** Make  $r$  the subject of the following formulae.

<b>a</b> $A = \pi r^2$	<b>b</b> $V = \frac{4}{3}\pi r^3$	<b>c</b> $P = \pi r + 2r$	<b>d</b> $V = \frac{2}{3}\pi r^2 h$
------------------------	-----------------------------------	---------------------------	-------------------------------------

**14** Make  $x$  the subject of the following formulae.

<b>a</b> $\frac{xy}{z} = \frac{ab}{cd}$	<b>b</b> $\frac{4\pi cx}{d} = \frac{3z}{py^2}$
---	--

**15** Make  $\sin B$  the subject of the formula  $\frac{a}{\sin A} = \frac{b}{\sin B}$

**16** Make  $\cos B$  the subject of the formula  $b^2 = a^2 + c^2 - 2ac \cos B$ .

## Extend

**17** Make  $x$  the subject of the following equations.

<b>a</b> $\frac{p}{q}(sx+t) = x-1$	<b>b</b> $\frac{p}{q}(ax+2y) = \frac{3p}{q^2}(x-y)$
------------------------------------	---



## Answers

$$1 \quad d = \frac{C}{\pi}$$

$$2 \quad w = \frac{P-2l}{2}$$

$$3 \quad T = \frac{S}{D}$$

$$4 \quad t = \frac{q-r}{p}$$

$$5 \quad t = \frac{2u}{2a-1}$$

$$6 \quad x = \frac{V}{a+4}$$

$$7 \quad y = 2 + 3x$$

$$8 \quad a = \frac{3x+1}{x+2}$$

$$9 \quad d = \frac{b-c}{x}$$

$$10 \quad g = \frac{2h+9}{7-h}$$

$$11 \quad e = \frac{1}{x+7}$$

$$12 \quad x = \frac{4y-3}{2+y}$$

$$13 \quad \mathbf{a} \quad r = \sqrt{\frac{A}{\pi}}$$

$$\mathbf{b} \quad r = \sqrt[3]{\frac{3V}{4\pi}}$$

$$\mathbf{c} \quad r = \frac{P}{\pi+2}$$

$$\mathbf{d} \quad r = \sqrt{\frac{3V}{2\pi h}}$$

$$14 \quad \mathbf{a} \quad x = \frac{abz}{cdy}$$

$$\mathbf{b} \quad x = \frac{3dz}{4\pi cpy^2}$$

$$15 \quad \sin B = \frac{b \sin A}{a}$$

$$16 \quad \cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$17 \quad \mathbf{a} \quad x = \frac{q+pt}{q-ps}$$

$$\mathbf{b} \quad x = \frac{3py+2pqy}{3p-apq} = \frac{y(3+2q)}{3-aq}$$